


CVS Chorionic Villus Sampling

T

A

A

G



You are being asked to consider prenatal diagnosis in your current pregnancy. Chorionic villus sampling (CVS) is available as a method of prenatal testing for women who are less than 13 weeks pregnant. This test can be performed earlier than amniocentesis, which is usually performed between 15 and 20 weeks, thereby offering earlier results. This brochure is to provide you with some basic information on CVS. A separate brochure is available which describes amniocentesis. A separate brochure is available which describes amniocentesis.

Who should consider CVS?

CVS should be considered by women age 35 or older at the time of delivery, individuals who have had a child with a chromosome abnormality, individuals who have a chromosome translocation, and couples at risk for a prenatally diagnosable genetic disease (e.g., hemophilia or sickle cell disease). CVS is not appropriate for individuals with a family history of neural tube defects (spina bifida or anencephaly).

When is CVS performed?

CVS is traditionally performed between 10 - 13 weeks after a woman's last menstrual period (during the first trimester).

How is CVS performed?

There are two methods for obtaining chorionic villi. For many women, either method can be safely performed. First, an ultrasound evaluation is performed to locate the developing placenta and to date the pregnancy. Often, the placental location determines which method of CVS is more appropriate.

There are certain other obstetrical considerations which may make one method preferable, including uterine anatomy and vaginal infections.

TRANSCERVICAL CVS: A thin catheter (hollow tube) is inserted through the vagina and cervix. Ultrasound is used to guide the catheter to the edge of the developing placenta. A small amount of placental tissue is removed for analysis.

TRANSABDOMINAL CVS: This procedure is similar to amniocentesis. A site is identified on the woman's abdomen and with ultrasound guidance a needle is inserted, through the abdomen and placed at the placenta. A small amount of tissue is removed for analysis.

What are the risks of CVS?

Most miscarriages occur early in pregnancy regardless of whether any invasive testing has been performed. A test like an amniocentesis carries a risk for complications, including miscarriage, of about 1 in 200 (0.5%). The risk after a CVS procedure is slightly higher (about 1%) To help keep the risk as low as possible, women are given precautionary instructions after the procedure. In 1991, there were several reports of babies with missing or shortened fingers or toes born to women who had undergone CVS. These birth defects were noted mainly when CVS was performed before the tenth week of pregnancy. As a result, CVS is now performed only during or after the tenth week of pregnancy and all women being offered CVS are informed about the possible small risk of limb defects. Studies on the risk of limb defects following CVS have had conflicting results. Some studies have found no increased risk, while others suggest that CVS may cause these limb abnormalities in about 1 in 3,000 births. However, recent studies suggest that CVS after 10 weeks of pregnancy poses a small or no increased risk for these limb defects. A study by the World Health Organization in 1999 found no increased risk for these abnormalities in babies born to more than 200,000 women who had CVS.

Based upon currently available information, the following can be concluded:

- 1) CVS is a relatively safe and accurate procedure and is considered an acceptable alternative to genetic amniocentesis.
- 2) CVS is best performed at 10–13 weeks pregnancy based on the beginning of the last menstrual period.
- 3) Although it is uncertain whether there is an increased risk of limb abnormalities following CVS, it is important to understand that this controversy has been raised and that the absolute risk may be in the order of 1 in 3,000 births. It is also important to remember that 2-3% of all children are born with a birth defect regardless of whether or not a woman had prenatal testing during her pregnancy.

What is analyzed from CVS?

The chorion is the outer membrane of the sac which surrounds the fetus. Early in pregnancy, the chorion is covered by finger-like projections called villi. In one area, the villi will implant in the uterus to develop into the placenta and the remainder of the villi disappear. The chorionic villi develop from the same fertilized egg as the fetus so the genetic material should be the same. The cells from the villi are cultured (grown) in the laboratory. Chromosome analysis and other special genetic testing, when indicated, can be performed on these cells.

What can be detected through CVS?

CVS detects chromosome abnormalities, such as the kind that causes Down syndrome. If indicated, some special genetic diseases can be diagnosed through DNA or enzyme analysis. CVS cannot detect neural tube defects such as spina bifida. Therefore, it is recommended that all women who undergo CVS consider having their blood drawn at 15-18 weeks of pregnancy for a spina bifida screening test (maternal serum AFP assay). This

can be arranged through your obstetrician. Many women also wish to pursue a level II ultrasound at 18-20 weeks for further information regarding the physical development of the fetus. No method of prenatal testing (CVS or amniocentesis) can guarantee a baby will be born without birth defects or genetic disease. Prenatal testing can only identify the diagnosable problems for which a couple is known to be at risk. Unfortunately, there are many conditions for which prenatal diagnosis is not yet available.

How accurate are the results from CVS?

Chromosome results from CVS are greater than 99% accurate. Occasionally, these results need to be clarified through blood tests on the parents, ultrasounds, amniocentesis, or fetal blood sampling.

When will I receive my test results?

Your CVS results will be available approximately 2 weeks after the procedure. Special studies for other genetic disease may take longer.

What are the advantages of CVS?

The greatest advantage of CVS is timing. Since CVS is performed during the first trimester of pregnancy, the results from the CVS are available earlier (12-15 weeks) than are amniocentesis results (17-22 weeks).

How Do I Learn More?

We hope this brochure answers some of your questions about CVS. All the information will be discussed in greater detail during the genetic counseling session, thus it is helpful to schedule an appointment for genetic counseling prior to the CVS procedure. The genetic counselor will not only discuss the procedure in greater detail, but will also determine whether additional testing is appropriate and ensure that all of your concerns are addressed. The genetic counselor is also available by phone or in person to follow-up with you after the results of the CVS become available.



Medical Genetics Laboratories

One Baylor Plaza, Nab 2015, Houston, Texas 77030

713-798-6555 • 1-800-411-gene (4363)

www.bcmgeneticlabs.org

BCM
Baylor College of Medicine