



March 2006

Vol. 11 No. 3

Rhinovirus was the most common virus isolated in February! Influenza A and B virus and RSV activity remained strong in February, and is persisting during the first half of March. Watch the RSV-FLU-O-METER

[http://www.bcm.edu/pedi/infect/dvl/newsletter/RSV-FLU-O-METER\(combined\).pdf](http://www.bcm.edu/pedi/infect/dvl/newsletter/RSV-FLU-O-METER(combined).pdf)

and "What's going 'round now!"

<http://www.bcm.edu/pedi/infect/dvl/updates.htm>.

Gail J. Demmler, M.D., Director
Diagnostic Virology Laboratory
Texas Children's Hospital
6621 Fannin Street, MC A370.17
3rd Floor, Room A 0370.01
Houston, TX 77030-2399, U.S.A.

(832) 824-2280 Telephone

(832) 825-2283 Fax



gjdemmler@texaschildrenshospital.org

EIA Rapid Test Results – February 2006

	RSV	FLU A	FLU B
Tested	609	668	668
Positive	37	48	8
% Positive	6	7	1

Viral Cultures – February 2006

Total all	946	Total Resp.	780
Positive	173	Positive	156
% Positive	18	% Positive	20



Visit our website for weekly updates @: www.bcm.tmc.edu/pedi/infect/dvl

<<http://www.texaschildrenshospital.org/Web/progressnotes/default.html>>