

Spinal Dural Arteriovenous Fistula and Concomitant Intramedullary Spinal Lesion


Fábio A. Nascimento¹, Peter Kan², Lydia Sharp¹, Jacob J. Mandel¹

¹Department of Neurology, Baylor College of Medicine, Houston, Texas, USA. ²Department of Neurosurgery, Baylor College of Medicine, Houston, Texas, USA.

CASE REPORT

We report a 63-year-old man who presented with a 6-month history of lower back pain followed by progressive bilateral leg weakness/paresthesia as well as bowel and bladder incontinence. Examination revealed proximal bilateral lower extremity weakness (3/5) and bilateral lower extremity sensory deficits. Upper extremity strength and sensation were normal. Reflexes were normal in all extremities; Babinski was absent. Patient was unable to walk due to lower extremity weakness.

Neuroimaging showed an enhancing C6 intramedullary lesion and diffuse T2 signal abnormality in the thoracic cord (extending from T5-T6 through the conus medullaris) as well as vascular flow voids along the dorsal surface of the lower cord to the conus medullaris (**figure 1A-1D**). Cerebrospinal fluid examination was unremarkable. He received a 5-day course of IV steroids, with minimal symptomatic improvement. Further investigation with spinal angiography revealed a L1 dural arteriovenous fistula (dAVF) (**figure 1E**). The dAVF was subsequently embolized with significant clinical improvement (**figure 1F**). Three-month follow-up spinal/cerebral angiography revealed no evidence of any vascular malformation in the cervical topography. Six-month follow-up imaging showed that the intramedullary cervical lesion had remained stable.



Spinal MRI: sagittal T2 (**A**) and postcontrast sagittal T1 (**B**) show an enhancing C6 9mm intramedullary lesion with mild expansion of the cord and focal edema; (**C**) sagittal T2 and postcontrast sagittal T1 (**D**) show diffuse signal abnormality from T5-T6 to the conus medullaris and vascular flow voids along the dorsal surface of the lower cord to the conus medullaris. **Spinal angiography (E,F).** Pre-embolization spinal angiography (**E**): anteroposterior injection of the left L1 segmental artery shows a type 1 dural arteriovenous fistula; note arterialized vein at midline. Post-embolization spinal angiography (**F**) shows Onyx in the left L1 segmental artery (previous point of fistularization) and complete obliteration of the arterialized vein without residual shunting.

DISCUSSION

Spinal dAVFs are the most common spinal vascular malformations, accounting for ~70% of such lesions. MRI findings suggestive of dAVFs include (a) centromedullary T2 hyperintensity over multiple segments, (b) T2 flow voids, often more pronounced on the dorsal surface of the cord and (c) diffuse T1 contrast enhancement within the cord.¹

Occurrence of a spinal dAVF and concurrent intramedullary spinal neoplastic lesion has not yet been described. Regarding our patient, although we cannot provide a definite diagnosis for his cervical lesion, we believe it is highly suspicious for low-grade glioma. Therefore, this patient may be the first reported case of spinal dAVF and concomitant intramedullary spinal neoplasm.

Dural arteriovenous fistula is a rare but treatable cause of progressive paraparesis. This condition should be included in the differential diagnoses of patients (especially older men) presenting with progressive symptoms of myelopathy whose imaging studies show an enlarged cord with intramedullary enhancement and T2 hyperintensity along with flow voids over the dorsal aspect of the cord. This report highlights the importance of pursuing a diagnosis of dAVF even in the setting of a concomitant, non-vascular spinal lesion.^{1,2,3}

DOWNLOAD POSTER / CONTACT INFORMATION



Dr. Fábio A. Nascimento
Baylor College of Medicine
7200 Cambridge Street, 9th floor | Houston, TX, 77030 | USA
Email: nascimento.fabio.a@gmail.com

REFERENCES

1. Krings T, Geibprasert S. Spinal dural arteriovenous fistulas. *AJNR Am J Neuroradiol.* 2009;30(4):639-648.
2. Koenig E, Thron A, Schrader V, Dichgans J. Spinal arteriovenous malformations and fistulae: clinical, neuroradiological and neurophysiologic findings. *J Neurol.* 1989;236(5):260-266.
3. Kim DJ, Willinsky R, Geibprasert S, et al. Angiographic characteristics and treatment of cervical spinal dural arteriovenous shunts. *AJNR Am J Neuroradiol.* 2010;31(8):1512-1515.
4. Morris JM. Imaging of dural arteriovenous fistula. *Radiol Clin North Am.* 2012;50(4):823-839.