



A Randomized, Double-Blind, Placebo-Controlled Study of Atomoxetine for Freezing of Gait in Parkinson's Disease

Muhammad M. Nashatizadeh, M.D., J. Ernesto Jimenez, M.Ed., Anthony L. Davidson, B.S., and Joseph Jankovic, M.D.

Parkinson's Disease Center and Movement Disorders Clinic

Department of Neurology, Baylor College of Medicine, Houston, Texas



OBJECTIVE

To evaluate the benefit of atomoxetine on freezing of gait (FoG) in Parkinson's disease (PD).

BACKGROUND

FoG is a common symptom in up to one-third of patients with longstanding or advanced PD. Various medications, surgical options and behavioral therapies have been proposed but patients often respond poorly or demonstrate inconsistent outcomes. Because a noradrenergic deficiency has been postulated to play a role in FoG, we designed a pilot study of the selective norepinephrine reuptake inhibitor atomoxetine in FoG related to PD.

DESIGN / METHODS

Five patients with FoG in PD were randomized to receive either active atomoxetine or placebo. All evaluators were blinded to treatment. Those receiving active treatment began on atomoxetine 10mg daily and the dose was escalated by 10mg increments up to 40mg over three weeks. Participants were evaluated at screening to verify eligibility, then at baseline and at two 4-week intervals, then after a two-week washout period. Scales administered were the Unified Parkinson's Disease Rating Scale (UPDRS), Gait and Balance Scale (GABS), and Clinician's Global Index of Change (CGIC). The subjects were also asked to complete the FoG Questionnaire (FOGQ) for evaluation of subjective improvement. Patients were videotaped performing tasks associated with the seven-meter step time (7MS) and the videos were rated by a blinded rater.

RESULTS

Three male patients and two female patients participated. Mean age was 65.6 ± 10.1 years. All were classified as either stage 3 or 4 on the Hoehn and Yahr (H&Y) scale and scored between 70-90 on the Schwab and England Activities of Daily Living (S-E ADL) scale. Three received active drug (patients 2, 4 and 5) while two received placebo (patients 1 and 3). No consistent differences in UPDRS Part I, II, or III scores were noted before or after treatment with either drug or placebo including question 14 (not shown) regarding subjective freezing. No consistent changes were noted in step number or duration on the 7MS test, in subjective improvement on the FOGQ, in GABS subscale or total scores, or in CGIC (not shown).

CONCLUSIONS

Although some patients reported subjective improvement, no consistent changes were demonstrated between atomoxetine and placebo in a small sample of PD patients with FoG. Further studies, using a large sample of subjects, may be needed to demonstrate atomoxetine's efficacy in the treatment of FoG.

TABLE 1: Patient Characteristics

Patient	Age/Sex	Diagnosed (yrs)	H&Y	S-E ADL
1	58/M	5.2	4	70
2	70/M	3.6	3	80
3	68/M	5.7	4	90
4	52/F	4.5	3	80
5	78/F	19.1	3	90

TABLE 2: Changes in GABS Scores

Patient	Subtotal I		Subtotal II		Total I+II		Fast Walk (sec)		Timed (sec)	
	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose
1	14	16	22	26	36	42	7	6	17	20
2	11	9	13	8	24	17	6	7	15	15
3	13	16	12	12	25	28	4	4	11	10
4	19	10	20	29	39	39	7	7	33	28
5	23	22	16	18	39	40	6	7	16	18

TABLE 3: Changes in 7MS Scores

Patient	Duration (sec)		Total Steps		FoG Episodes	
	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose
1	25	24	37	44	1	0
2	23	20	38	32	0	0
3	14	19	24	34	0	1
4	24	22	53	47	0	1
5	19	25	38	46	0	0

TABLE 4: Changes in UPDRS Scores

Patient	UPDRS Part I		UPDRS Part II		UPDRS Part III	
	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose
1	6	6	24	25	38	28
2	4	5	18	20	30	26
3	2	3	23	22	42	35
4	1	3	17	9	45	48
5	4	4	25	26	47	47

TABLE 5: Changes in FoGQ Scores

Patient	Worst State		ADL Difficulty		Glued to Floor		Longest Freezing		Start Hesitation		Turn Hesitation		Total	
	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose	Baseline	Max Dose
1	3	2	2	2	4	4	3	2	2	2	2	2	16	15
2	3	3	2.5	2	3.5	4	2	2	1	2	1	1	13	14
3	3	3	4	3	4	3	4	3	4	3	3	3	22	18
4	3	2	2	1	3	2	2	2	3	2	2	2	15	11
5	3	3	3	3	3	3	2	2	2	1	1	0	13	12

REFERENCES

1. Amboni M, Cozzolino A, Longo K et al. Freezing of gait and executive functions in patients with Parkinson's disease. *Mov Disord*. 2008 Feb 15;23(3):395-400.
2. Moore O, Perez C, Giladi N. Freezing of gait affects quality of life of peoples with Parkinson's disease beyond its relationships with mobility and gait. *Mov Disord*. 2007 Nov 15;22(15):2192-5.
3. Tinetti ME, Williams TF, Mayewski R. Fall risk index for elderly patients based on number of chronic disabilities. *Am J Med*. 1986;80:429-434.
4. Fahn S, Marsden CD, Calne DB, Goldstein M, eds. Recent Developments in Parkinson's Disease, Vol. 2. Florham Park, NJ: Macmillan Health Care Information 1987, pp 153-163, 293-304.
5. Thomas M, Jankovic J, Suteerawattananon M et al. Clinical gait and balance scale (GABS): validation and utilization. *J Neurol Sci*. 2004 Jun 15;221(1-2):125.
6. Giladi N, Shabtai H, Simon ES et al. Construction of freezing of gait questionnaire for patients with Parkinsonism. *Parkinsonism and Related Disord* 2000;6:165-170.
7. Guy W and Bonato RR, eds. Manual for the ECDEU Assessment Battery, 2nd Revised ed. Chevy Chase, MD: National Institute of Mental Health; 1970:12-1-12-6.
8. Webster, DD. Critical analysis of the disability in Parkinson's disease. *Modern Treatment* 1968;5(2):257-282.
9. Weber PC and Cass SP. Clinical assessment of postural stability. *Am J Otolary* 1993;14:566-569.
10. Giladi N, McMahon D, Przedborski S et al. Motor blocks in Parkinson's disease. *Neurology* 1992;42:333-339.
11. Corboy DL, Wagner ML and Sage JI. Apomorphine for motor fluctuations and freezing in Parkinson's disease. *Ann Pharmacother* 1995;29:282-288.
12. Ogawa N, Yamamoto M and Takayama H. L-threo-3,4-dihydroxyphenylserine treatment of Parkinson's disease. *J Med* 1985;16:525-534.
13. Tohgi H, Abe T, Takahashi S et al. Effect of a synthetic norepinephrine precursor, L-threo-3,4-dihydroxyphenylserine on the total norepinephrine concentration in the cerebrospinal fluid of parkinsonian patients. *Neurosci Lett* 1990;116:194-197.
14. Tohgi H, Abe T, and Takahashi S. The effects of L-threo-3,4-dihydroxyphenylserine on the total norepinephrine and dopamine concentrations in the cerebrospinal fluid and freezing gait in parkinsonian patients. *J Neural Transm Park Dis Dem Sect* 1993;5:27-34.
15. Katsube J, Narabayashi H, Hayashi A et al. [Development of L-threo-DOPS, a norepinephrine precursor amino acids]. *Yakugaku Zasshi* 1994;114:823-846.

This research study was funded by Eli Lilly and Company.