

Transcription Factor PITX3 Gene in Parkinson's Disease

Weidong Le^{*}, a, Deanne Nguyen^a, Maosheng Huang^b, Yujian Ding^c, Hao Deng^d, Joseph Jankovic^a



a. Department of Neurology, Baylor College, Houston, TX, USA

b. Department of Epidemiology, MD. Anderson Cancer Center, Houston, TX, USA

c. Institute of Neurology, Ruijin Hospital, Shanghai Jiao Tong University, Shanghai, China

d. Department of Neurology, the Third Xiangya Hospital, Changsha, China

Abstract

PITX3 is a transcription factor important for the differentiation and survival of midbrain dopaminergic neurons during the development. Single nucleotide polymorphisms (SNP) in the gene may be associated with Parkinson's Disease (PD). To verify their findings and to determine the nature of the association in a subset of our PD patients we have analyzed two *PITX3* SNPs (rs2281983 and rs4919621) in PD patients and age-matched health controls. Our data show that the substitutions of C/T in SNP1 and A/T in SNP2 are significantly higher in PD, and this finding is even more robust in young onset and familial PD as compared with age-matched healthy controls. Our findings indicate that *PITX3* may play a role in the pathogenesis of PD.

Material and Methods

Patients: 265 North American Caucasian PD patients and 210 healthy controls from the Parkinson's Disease Center and Movement Disorders Clinic (PDCMDC), Baylor College of Medicine (BCM). All research subjects signed an informed consent, approved by BCM institutional Review Board for Human Research. PD samples were subcategorized into EOPD, LOPD, familial PD and sporadic PD.

Data Table 1.

Allele	Controls	Total PD	EOPD	LOPD	fPD	sPD
N (allele)	420	530	162	368	196	334
Gender M/F	101/109	143/122	41/40	102/82	53/45	90/77
Age	57.92±13.87	59.18±12.62	44.91±9.08	64.86±14.17	54.74±15.92	62.24±12.80

PCR: Genomic DNA was extracted from peripheral blood using standard protocols. PCR products of *PITX3* were amplified with the following primers:

rs2281983	Forward	5'-CGGGTCTGAGAGCATACC-3'
	Reverse	5'-GACGGTTCGCTGAAAAGAA-3'
rs4919621	Forward	5'-CTGATCCTCCAAACCCCTGA-3'
	Reverse	5'-AGATGAGGGGGCCCTTTAGA-3'

Statistics: Pearson's χ^2 tests were applied to test for significance in differences of gene frequencies. A value ($p < 0.05$, two-tailed) was considered to be significant.

Results




Figure 1. The gene loci of SNP1 rs2281983 and SNP2 rs4919621

Allele	Controls	Total PD	EOPD	LOPD	fPD	sPD
rs2281983						
T	248 (0.590)	271 (0.511)	78 (0.481)	193 (0.524)	89 (0.454)	182 (0.545)
C	172 (0.410)	259 (0.489)	84 (0.519)	175 (0.476)	107 (0.546)	152 (0.455)
Empirical-P		0.035	0.038	0.131	0.005	0.291
Adjusted OR* (95%CI)	Reference	1.34 (1.03-1.74)	2.05 (1.33-3.16)	1.24 (0.92-1.68)	1.72 (1.22-2.43)	1.18 (0.88-1.59)

	p -Value ^a	p -Value ^b	p -Value ^c	p -Value ^d	p -Value ^e
C allele:	0.015	0.018	0.063	0.002	0.209

Data Table 2. Gender: male vs female (M/F);

Age: Mean \pm SD; Significant ($P < 0.05$)

P-Value a : All PD when compared to controls

P-Value b : EOPD when compared to controls

P-Value c : LOPD when compared to controls

P-Value d : fPD when compared to controls

P-Value e : sPD when compared to controls

*: adjusted by sex and age

Allele	Controls	Total PD	EOPD	LOPD	fPD	sPD
rs4919621						
T	268 (0.638)	294 (0.555)	87 (0.537)	207 (0.562)	103 (0.525)	191 (0.572)
A	152 (0.360)	236 (0.445)	75 (0.463)	161 (0.438)	93 (0.475)	143 (0.428)
Empirical-P		0.022	0.018	0.057	0.017	0.078
Adjusted OR (95%CI)	Reference	1.47 (1.07-1.92)	2.17 (1.41-3.36)	1.21 (0.89-1.64)	1.78 (1.07-2.58)	1.20 (0.96-1.53)

	p -Value ^a	p -Value ^b	p -Value ^c	p -Value ^d	p -Value ^e
A allele:	0.009	0.025	0.030	0.008	0.064

Data Table 3. Gender: male vs female (M/F)

Age: Mean \pm SD; Significant ($P < 0.05$)

P-Value a : All PD when compared to controls

P-Value b : EOPD when compared to controls

P-Value c : LOPD when compared to controls

P-Value d : fPD when compared to controls

P-Value e : sPD when compared to controls

*: adjusted by sex and age

Conclusions

1. The frequency of substitutions of C/T in SNP1 rs2281983 and A/T in SNP2 rs4919621 is significantly higher in PD, and it is particularly high in EOPD and fPD as compared with age-matched healthy controls.

2. Our results support the hypothesis that *PITX3* is critical not only in the development and maintenance of the dopaminergic system, but also in the pathogenesis of PD.

Acknowledgement

This study was supported by the Helis Foundation (W.L.) the National Parkinson Foundation to the Baylor College of Medicine Center of Excellence (J.J.) the National Natural Scientific Foundation of China (30871351, H.D.). Thank Dr. William Ondo, and Anthony L. Davidson for their efforts in collecting the genetic information and DNA specimens.

References

- Deng H, Le W.D, Zhang X, Pan T.H, Jankovic J. G309 and W437OPA PINK1 mutations in Caucasian Parkinson's disease patients. *Acta Neurol Scand* 111, 6:351-352.
- Bergman O, Hakansson A, Westberg L, Nordenstrom K, Nissbrandt H. PITX3 polymorphism is associated with early onset Parkinson's disease. *Neurobiol Aging* [Epub ahead of print]
- Peng C, Fan S, Li X, Fan X, Le W. Overexpression of pitx3 upregulates expression of BDNF and GDNF in SH-SY5Y cells and primary ventral mesencephalic cultures. *FEBS* 581, 7:1357-6.



The movement Disorder Society