





January 2008

Vol. 13 No. 1

RSV was the most common virus isolated, followed by influenza virus – both types A and B. Adenoviruses, rhinoviruses and parainfluenza viruses were also detected. Watch the RSV-FLU-O-METER for weekly updates

[http://www.bcm.edu/pedi/infect/dvl/newsletter/RSV-FLU-O-METER\(combined\).pdf](http://www.bcm.edu/pedi/infect/dvl/newsletter/RSV-FLU-O-METER(combined).pdf) and Check out "What's going 'round now!" updates <http://www.bcm.edu/pedi/infect/dvl/>



Gail J. Demmler, M.D., Director
Diagnostic Virology Laboratory

Texas Children's Hospital
6621 Fannin Street, MC A370.17
3rd Floor, Room AB 0370.01
Houston, TX 77030-2399, USA

832 824 2280 Telephone
832 825 2283 Fax
gidemmler@texaschildrenshospital.org

Viral Cultures – Dec 2007

Total all	1352	Total Resp.	1160
Positive	304	Positive	294
% Positive	22	% Positive	25

EIA Rapid Test Results – Dec 2007

	RSV	FLU A	FLU B	ADENO
Tested	943	1010	1010	226
Positive	234	61	4	2
% Positive	25	6	0	1